



In preparation for your upcoming surgery, please review the following reminders:

INSTRUCTIONS FOR YOUR LENS PROCEDURE

- **General Reminders:**
 - A member of our team will call you the day before your procedure to review your medical history and to let know your surgery appointment time.
 - Please fill your eyedrop prescriptions if you have not already done so.
- **Start using the following eyedrops 3 days prior to your procedure:**
 - Place 1 DROP of ILEVRO 1 TIME EACH DAY and 1 DROP of MOXIFLOXACIN (VIGAMOX) 3 TIMES EACH DAY (breakfast, lunch, and bedtime) in your operative eye(s).
 - Please wait 5 minutes between eye drops.
 - **DO NOT USE DUREZOL.** You will start using this drop **after** your procedure.
- **On the Morning of Your Procedure:**
 - Place 1 DROP of ILEVRO and MOXIFLOXACIN (VIGAMOX) in your operative eye(s).
 - Please wait 5 minutes between eye drops.
 - **DO NOT USE DUREZOL.** You will start using this drop **after** your procedure.
- **Additional Important Information:**
 - **DO NOT** eat or drink **ANYTHING** at least 8 HOURS prior to your surgery. No gum or mints. You may brush your teeth, but do not swallow water.
 - You may take your regular medications in the morning with a sip of water (AVOID diabetic medications or diuretics).
 - **DO NOT** apply cologne or perfume.
 - **DO NOT** apply makeup or lotions to your face.
 - Wear loose comfortable clothing: preferably a shirt or blouse that opens in the front. Leave jewelry and other valuables at home.
 - HAVE A RESPONSIBLE ADULT ACCOMPANY AND DRIVE YOU TO THE CENTER AND BACK HOME.
 - Notify the office of any change in your health (fever, cold, sore throat, etc.) before the procedure.

If you have any additional questions, please call SightTrust at (954) 653-0100.

Prescription Substitutions

Due to the recent prescription's shortages SightTrust has approved a list of substitutions:

Substitute for Vigamox

Moxifloxacin

Instructions for use:

1 drop 3 times per day for 1 week

Substitute for Ilevro

Prolensa

Instructions for use:

1 drop 1 time per day for 6 weeks

Substitute for Durezol (for use after surgery)

Difluprednate

Instructions for use:

1 drop 2 times per day for 2 weeks,
then 1 drop 1 time per day for 2 weeks,

If you have any additional questions, please call SightTrust at (954) 653-0100.